

The ATLAS Diboson Resonance in Non-Supersymmetric SO(10)

Jason L. Evans^{1,2}, Natsumi Nagata², Keith A. Olive², and Jiaming Zheng²

¹*School of Physics, KIAS, Seoul 130-722, Korea*

²*William I. Fine Theoretical Physics Institute, School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

Abstract

SO(10) grand unification accommodates intermediate gauge symmetries with which gauge coupling unification can be realized without supersymmetry. In this paper, we discuss the possibility that a new massive gauge boson associated with an intermediate gauge symmetry explains the excess observed in the diboson resonance search recently reported by the ATLAS experiment. The model we find has two intermediate symmetries, $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ and $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$, where the latter gauge group is broken at the TeV scale. This model achieves gauge coupling unification with a unification scale sufficiently high to avoid proton decay. In addition, this model provides a good dark matter candidates, whose stability is guaranteed by a $\mathbb{Z}_2$ symmetry present after the spontaneous breaking of the intermediate gauge symmetries. We also discuss prospects for testing these models in the forthcoming LHC experiments and dark matter detection experiments.

1 Introduction

Grand unification [1] is considered to be one of the most promising frameworks for physics beyond the Standard Model (SM). SO(10) Grand Unified Theories (GUTs) [2–4] have many especially attractive features. First, the full set of SM fermions of each generation together with a right-handed neutrino is embedded into a single **16** chiral representation of SO(10). Second, anomaly cancellation in the SM can be understood naturally since SO(10) is free from anomalies. Third, gauge coupling unification can be realized without relying on supersymmetry, if there are intermediate-scale gauge symmetries [5]. Majorana mass terms of right-handed neutrinos can be generated at an intermediate scale, which may explain the smallness of neutrino masses. These interesting aspects have stimulated many extensive studies of SO(10) GUTs [6–9].

In addition, SO(10) GUTs with intermediate gauge symmetries predict new gauge bosons whose masses are of the order of the breaking scales of the corresponding gauge symmetries. The breaking scale of these intermediate gauge symmetries affects the running of the gauge couplings. These scales can be determined by requiring gauge coupling unification, which is highly dependent on the symmetry-breaking pattern and matter content of the model. However, we will show that it is possible to obtain a model containing TeV-scale gauge bosons accessible at the LHC, for a particular choice of the intermediate gauge symmetries and particle content.

In this paper, we consider non-supersymmetric SO(10) GUT models with an intermediate gauge symmetry showing up around the TeV scale, and discuss the possibility that a new gauge boson associated with the intermediate gauge symmetry can explain the anomalous diboson events recently reported by the ATLAS collaboration [10]. We will discuss a model with two intermediate symmetries, $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ which is then broken to $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$. The unified and intermediate scales are determined by the renormalization group (RG) running of the gauge couplings and depend on the matter content of the theory. In the model discussed, the GUT scale is of order 10^{17} GeV and is high enough so that the proton decay bounds can be evaded. The first intermediate scale is of order 10^9 GeV and is high enough that constraints from leptoquarks are satisfied. The second intermediate scale lies around a few TeV and the charged gauge bosons of $SU(2)_R$, $W_R^\pm$, can have a mass ~ 2 TeV and may explain the excess observed in the ATLAS diboson resonance search [10]. The possibility of this new charged gauge boson explaining the diboson anomaly is discussed in Refs. [11–16]. We will see that $W_R^\pm$ in this model can also reproduce the diboson excess. Furthermore, this $W_R^\pm$ may explain another anomaly observed by the CMS collaboration in the right-handed neutrino searches with a dijet and a dilepton final state [17], as we will discuss below. Our model also predicts an additional massive boson: a neutral gauge boson Z_R . Detection of this Z_R would be a smoking-gun signature of our model. The masses of the Z_R can be heavier than the $W_R^\pm$ mass, and thus evades the current LHC bound. However, it may be in reach of Run-II of the LHC, as we will discuss.

Our model also contains a promising dark matter candidate. The dark matter candidate is stabilized by a residual $\mathbb{Z}_2$ symmetry arising after the SO(10) symmetry has

been completely broken to the SM gauge symmetries [18–27]. This residual $\mathbb{Z}_2$, as it turns out, is equivalent to matter parity [28]. It is tantalizing that the presence of this dark matter candidate is essential for the model to achieve gauge coupling unification. An integral part of obtaining gauge coupling unification in this set up is an intermediate scale of order a TeV, a second intermediate scale large enough to suppress the effects of leptoquarks, and a GUT scale which is large enough to avoid proton decay.¹ Given these conditions, the required matter content at low energies is severely restricted, and thus it is highly non-trivial that our model contains a good dark matter candidate. The presence of dark matter gives us another opportunity to test this model; we can probe the dark matter candidate predicted in this model in dark matter detection experiments. We also discuss this prospect in what follows. Combining dark matter search results with those obtained at the LHC experiments, we may be able to test our model in near future.

This paper is organized as follows. We first describe our model in the next section. Then, in Sec. 3, we briefly review the ATLAS diboson anomaly, and show that the $W_R^\pm$ in our model can explain the diboson anomaly. We also discuss the present constraints on our models from various experiments and consider the testability of this model in future LHC experiments and dark matter searches. Finally, we conclude in Sec. 4.

2 The Model

In this section, we describe the specific SO(10) model used in this analysis. This model is obtained in a similar manner to that discussed in Ref. [25], except now we allow for two intermediate scales. To obtain these intermediate scales, we need to arrange some Higgs fields to acquire vacuum expectation values (VEVs) in specific directions at particular scales. Throughout this work, we make an assumption on the Higgs field content. We assume that at each intermediate scale the Higgs fields present are components of the SO(10) multiplets that have VEVs equal to or less than the corresponding intermediate scale. The other components have masses of order of the previous symmetry-breaking scale. We also allow for a light SO(10) component which could be dark matter.

Before we get into the details of our model, we first discuss the necessity for two intermediate scales. If there were only one intermediate scale and we wished to explain the ATLAS diboson excess, we would have to choose the TeV scale. For intermediate gauge symmetries which include the $SU(4)_C$ subgroup, this choice turns out to be problematic since the breaking of the $SU(4)_C$ subgroup generates a vector leptoquark at that intermediate scale. Since this leptoquark interacts with both left- and right-handed fermions, it will generate excessively large contributions to the meson decays (see, *e.g.*, Ref. [32]). In fact, one can show that no tuning can simultaneously tune away the contribution of a vector leptoquark to the Kaon decays $K^+ \rightarrow \pi^+ \bar{e} \mu$ and $K_L \rightarrow \bar{e} \mu$ by exploiting the contributions of the scalar leptoquarks contained inside the Higgs fields.² If the interme-

¹Realization of a few TeV intermediate scale within the framework of grand unification is also discussed in Refs. [29–31].

²This is because both of these decays arise from the same effective operator. These two decays depend

mediate gauge symmetries do not contain the $SU(4)_C$ subgroup, on the other hand, we find that we cannot obtain gauge coupling unification with a sufficiently high GUT scale since there are no new contributions to the running of the $SU(3)_C$ gauge coupling constant. This makes it impossible to get the intermediate scale to be around the TeV scale while maintaining gauge coupling unification with a sufficiently high GUT scale. These problems, however, can be evaded if a two step breaking pattern is considered. The vector leptoquark then has a mass of order of the larger intermediate scale. This larger intermediate scale generates a mass for the vector leptoquarks which is high enough to avoid constraints coming from meson decays but low enough that the vector leptoquarks still contribute significantly to the running of the $SU(3)_C$ gauge coupling.

The breaking pattern of our model, with two intermediate gauge symmetries, is from $SO(10)$ to $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ which is then broken to $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$. For the breaking pattern we consider, $SO(10)$ is broken by the VEV of the $(\mathbf{1}, \mathbf{1}, \mathbf{1})$ component of a $\mathbf{210}$ into $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$. This symmetry is then subsequently broken by a VEV of the $(\mathbf{15}, \mathbf{1}, \mathbf{1})$ component of the same $\mathbf{210}$ to $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$. Importantly, the $(\mathbf{1}, \mathbf{1}, \mathbf{1})$ component of the $\mathbf{210}$ breaks D -parity [33] at the GUT scale, and thus the low-energy effective theory does not exhibit a left-right symmetry [34]. Finally, this gauge group is broken to the SM gauge symmetries at the TeV scale by the $(\mathbf{1}, \mathbf{1}, \mathbf{3}, +2)$ component of a $\mathbf{126}$. However, there remains a $\mathbb{Z}_2$ symmetry coming from $U(1)_{B-L}$ since the $B-L$ charge of this component of the $\mathbf{126}$ is 2 [18–21]. The $(\mathbf{1}, \mathbf{1}, \mathbf{3}, 0)$ component of the $\mathbf{210}$ also breaks $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$, but does not break the $\mathbb{Z}_2$ symmetry since its $B-L$ charge is zero. In summary, we consider the following symmetry-breaking chain:

$$\begin{aligned} SO(10) &\xrightarrow{\langle \mathbf{210} \rangle} SU(4)_C \otimes SU(2)_L \otimes SU(2)_R \\ &\xrightarrow{\langle \mathbf{210} \rangle} SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L} \\ &\xrightarrow{\langle \mathbf{126} \rangle \langle \mathbf{210} \rangle} SU(3)_C \otimes SU(2)_L \otimes U(1)_Y \otimes \mathbb{Z}_2 . \end{aligned} \quad (1)$$

Under the remnant $\mathbb{Z}_2$ symmetry, which is found to be equivalent to matter parity $P_M = (-1)^{3(B-L)}$ [28], the SM fermions are odd while the SM Higgs is even. Therefore, a scalar boson (fermion) can be stable if it is odd (even) under this $\mathbb{Z}_2$ symmetry. If such a particle is electrically and color neutral, it can be a good dark matter candidate [22, 23]. For a detailed discussions on this class of dark matter candidates in $SO(10)$ GUTs, see Refs. [24, 25].

The low-energy matter content of this model, beyond the SM fermions and right-handed neutrinos (which are embedded into three $\mathbf{16}$ representations), is given in Table 1. Here, the first, second, and third columns show the $SO(10)$, $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$, and $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ quantum numbers of the particles, respectively,

on the vector current and axial current respectively. Because the effective operators generated from scalar leptoquarks will have different chirality for the quarks than the vector leptoquark, the contribution from the scalar leptoquark will add to one decay and subtract from the others. Therefore, no tuning can remove this contribution.

while the fourth column represents their $\mathbb{Z}_2$ charges. The subscripts C and D refer to complex and Dirac respectively. Most of the components of the **210** have GUT scale masses and do not affect our analysis. In this model, the $(\mathbf{15}, \mathbf{1}, \mathbf{1})_C$ component of **210**_C has a mass similar to the $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ breaking scale, as its VEV breaks the symmetry. Similarly, the components of $(\mathbf{15}, \mathbf{1}, \mathbf{3}) \in \mathbf{210}$ and $(\overline{\mathbf{10}}, \mathbf{1}, \mathbf{3}) \in \mathbf{126}$ charged under color have masses of order the higher intermediate scale breaking. The components in the third column of the table are those components which are fine-tuned to have masses lighter than this scale. Among the light fields are the $(\mathbf{1}, \mathbf{1}, \mathbf{3}, +2)_C$ and the $(\mathbf{1}, \mathbf{1}, \mathbf{3}, 0)_C$ which obtain VEVs and break the $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ symmetry to the SM symmetries. The components of these fields which are not eaten by the gauge bosons receive masses of order of the VEV breaking this symmetry. The rest of the Higgs fields listed in the table are responsible for breaking the electroweak symmetry and have masses of order the electroweak scale. Also listed in the table is an $SU(2)_L$ triplet Dirac fermion which can be a viable dark matter candidate and could originate from a **45** of $SO(10)$. The remaining components of the **45** (beyond the triplet) should all have GUT scale masses. In summary, the states in column 2 affect the running of the gauge couplings between the high intermediate scale and the GUT scale, and those in column 3 affect the running between the TeV scale and the high intermediate scale. The particles in the third column and green shaded rows contribute to the running of the gauge couplings below the TeV scale. This leads to a theory below the TeV scale with the SM fermions and four doublet Higgs bosons³. A similar decomposition can be made for the gauge bosons; the $(\mathbf{6}, \mathbf{2}, \mathbf{2})$ component of the $SO(10)$ gauge boson, **45**, is around the GUT scale, while the other components, the $SU(4)_C$, $SU(2)_L$, and $SU(2)_R$ gauge bosons, are massless at this stage of the GUT symmetry breaking. A part of the $SU(4)_C$ gauge bosons, the vector leptoquark, obtains a mass of the order of the higher intermediate scale, as we will see below.

The mass of the dark matter candidate is fixed by requiring that its thermal relic abundance agree with the observed dark matter density $\Omega_{\text{DM}} h^2 \simeq 0.12$ [35]. The thermal relic abundance of any DM candidate is determined by its thermally averaged annihilation cross section σ_{ann} times the relative velocity between the initial particles, v_{rel} , via the following relation:

$$\Omega_{\text{DM}} h^2 \simeq \frac{3 \times 10^{-27} \text{ cm}^3 \text{ s}^{-1}}{\langle \sigma_{\text{ann}} v_{\text{rel}} \rangle} . \quad (2)$$

Here, the brackets signify a thermal average. In the case of an $SU(2)_L$ triplet Dirac fermion dark matter candidate with zero hypercharge, the s -wave annihilation cross section is given by [36]

$$\langle \sigma_{\text{ann}} v_{\text{rel}} \rangle \simeq \frac{37 g_2^4}{192 \pi m_{\text{DM}}^2} , \quad (3)$$

where g_2 denotes the $SU(2)_L$ gauge coupling constant and m_{DM} is the dark matter mass. From Eqs. (2) and (3), we obtain $m_{\text{DM}} \simeq 2 \text{ TeV}$. However, it is known that the non-perturbative Sommerfeld effect significantly enhances the annihilation cross section [37],

³As we will see below, the dark matter candidate will need to be of order the TeV scale.

Table 1: *Particle content of our model. The first, second, and third columns show the $SO(10)$, $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$, and $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ quantum numbers of the particles, respectively, while the forth column represents their $\mathbb{Z}_2$ charges. The orange (green) shaded fields acquire VEVs of about a few TeV (the electroweak scale), and the blue shaded particle contains a dark matter candidate. The $(\mathbf{15}, \mathbf{1}, \mathbf{1})_C$ of the $\mathbf{210}_C$ lies around the $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ breaking scale, and none of its components appears below this scale.*

$SO(10)$	$SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$	$SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$	$\mathbb{Z}_2$
$\mathbf{210}_C$	$(\mathbf{15}, \mathbf{1}, \mathbf{1})_C$	–	+
$\mathbf{126}_C$	$(\overline{\mathbf{10}}, \mathbf{1}, \mathbf{3})_C$	$(\mathbf{1}, \mathbf{1}, \mathbf{3}, +2)_C$	+
$\mathbf{210}_C$	$(\mathbf{15}, \mathbf{1}, \mathbf{3})_C$	$(\mathbf{1}, \mathbf{1}, \mathbf{3}, 0)_C$	+
$\mathbf{10}_C$	$(\mathbf{1}, \mathbf{2}, \mathbf{2})_C$	$(\mathbf{1}, \mathbf{2}, \mathbf{2}, 0)_C$	+
$\mathbf{126}_C$	$(\mathbf{15}, \mathbf{2}, \mathbf{2})_C$	$(\mathbf{1}, \mathbf{2}, \mathbf{2}, 0)_C$	+
$\mathbf{45}_D$	$(\mathbf{1}, \mathbf{3}, \mathbf{1})_D$	$(\mathbf{1}, \mathbf{3}, \mathbf{1}, 0)_D$	+

which results in a smaller cross section, and thus larger dark matter mass, needed to reproduce the correct dark matter density. As far as we know, the thermal relic abundance of a triplet Dirac fermion has not been computed yet with the Sommerfeld enhancement effect included. However, we expect that the favored mass for a Dirac triplet is smaller than that for a Majorana triplet (which was found to be ~ 2.7 TeV [38]), since the thermally-averaged annihilation cross section of Dirac dark matter is a factor of two smaller than that of Majorana dark matter.

In Fig. 1, we show a plot of the running of the gauge coupling constants in this model. Here, we set the masses of the fields in the third column in Table 1 to be 1.9 TeV. In this figure, we show the running of the inverse of $\alpha_a \equiv g_a^2/(4\pi)$ ($a = 1, 2, 3$) with g_1 , g_2 , and g_3 corresponding to the $U(1)_Y$, $SU(2)_L$, and $SU(3)_C$ gauge coupling constants, respectively. The definitions of these gauge couplings in the $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ theory are given by

$$\frac{1}{\alpha_1} \equiv \frac{3}{5} \frac{1}{\alpha_{2R}} + \frac{2}{5} \frac{1}{\alpha_4}, \quad \alpha_2 \equiv \alpha_{2L}, \quad \alpha_3 \equiv \alpha_4, \quad (4)$$

where α_4 , α_{2L} , and α_{2R} represent the $SU(4)_C$, $SU(2)_L$, and $SU(2)_R$ gauge couplings, respectively, while those in the $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ theory are

$$\frac{1}{\alpha_1} \equiv \frac{3}{5} \frac{1}{\alpha_{2R}} + \frac{2}{5} \frac{1}{\alpha_{B-L}}, \quad \alpha_2 \equiv \alpha_{2L}, \quad (5)$$

where α_{B-L} is the $U(1)_{B-L}$ gauge coupling. Using the weak scale values of the gauge couplings as inputs and insisting on unification of the coupling constants at some high energy scale, allows us to use the three RGEs to solve for the higher intermediate scale,

the GUT scale, and the value of the the unified gauge coupling at the GUT scale (recall that we have fixed the value of the lower intermediate scale). From this figure, we find that the GUT scale in this model is 9.8×10^{16} GeV, which is high enough to avoid proton decay constraints. The $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ breaking scale is determined to be $\simeq 7.2 \times 10^8$ GeV. The $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ gauge coupling constants at 1.9 TeV are found to be

$$g_{B-L} \simeq 0.58 , \quad g_{2L} \simeq 0.64 , \quad g_{2R} \simeq 0.42 , \quad g_3 \simeq 1.03 . \quad (6)$$

The value of g_{2R} above will be important when we discuss the diboson anomaly.




Figure 1: *Running of gauge couplings in our model. Here, we set the masses of the fields in the third column in Table 1 to be 1.9 TeV.*

In this model, only $W_R^\pm$ and Z_R lie around the TeV scale, while the vector leptoquarks have masses of $\mathcal{O}(10^{8-9})$ GeV. The scalar leptoquarks arising from the $(\mathbf{15}, \mathbf{1}, \mathbf{1})_C$ and $(\mathbf{15}, \mathbf{1}, \mathbf{3})_C$ of the $\mathbf{210}_C$ and $(\bar{\mathbf{10}}, \mathbf{1}, \mathbf{3})_C$ and $(\mathbf{15}, \mathbf{2}, \mathbf{2})_C$ of the $\mathbf{126}_C$ also have a mass of order this scale. The masses of $W_R^\pm$ and Z_R are computed to be

$$\begin{aligned} m_{W_R}^2 &\simeq g_{2R}^2 [(v_R^{\mathbf{10}})^2 + (v_R^{\mathbf{15}})^2] , \\ m_{Z_R}^2 &\simeq 2 \left(g_{2R}^2 + \frac{3}{2} g_{B-L}^2 \right) (v_R^{\mathbf{10}})^2 , \end{aligned} \quad (7)$$

where

$$\langle (\mathbf{1}, \mathbf{1}, \mathbf{3}, +2)_C \rangle = T_- v_R^{\mathbf{10}} , \quad \langle (\mathbf{1}, \mathbf{1}, \mathbf{3}, 0)_C \rangle = T_3 v_R^{\mathbf{15}} . \quad (8)$$

Here, $T_a = \tau_a/2$ ($a = 1, 2, 3$) with τ_a the Pauli matrices and $T_- \equiv T_1 - iT_2$, and we have neglected the contribution of the electroweak-scale VEVs for simplicity. On the other

hand, the vector leptoquarks, whose quantum numbers under the $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ symmetry are $(\mathbf{3}, \mathbf{1}, +2/3) \oplus (\bar{\mathbf{3}}, \mathbf{1}, -2/3)$, acquire a mass of

$$m_{V_{LQ}}^2 \simeq \frac{2}{3} g_A^2 (v_{422}^{\mathbf{15}})^2, \quad (9)$$

with

$$\langle (\mathbf{15}, \mathbf{1}, \mathbf{1})_C \rangle = X_{B-L} v_{422}^{\mathbf{15}}, \quad (10)$$

where X_{B-L} is the $SU(4)_C$ generator corresponding to the $B-L$ charge, i.e. $\sqrt{\frac{3}{8}}(B-L)$.⁴ As the $(\mathbf{15}, \mathbf{1}, \mathbf{1})_C$ VEV $v_{422}^{\mathbf{15}}$ sets the $SU(4)_C \otimes SU(2)_L \otimes SU(2)_L$ breaking scale, $m_{V_{LQ}}$ is as high as $\sim 10^9$ GeV. The masses of the scalar leptoquarks are dependent on the couplings in the scalar potential; generically, they are also $\mathcal{O}(v_{422}^{\mathbf{15}})$.

Since the gauge couplings g_{B-L} and g_{2R} are determined, and given in Eq. (6), if we fix the $W_R^\pm$ mass, then we obtain the Z_R mass as a function of $v_R^{\mathbf{15}}$. We show this in Fig. 2 setting $m_{W_R} = 1.9$ TeV. It is found that the Z_R mass is relatively small; in particular, m_{Z_R} can be much smaller than that predicted in the minimal left-right symmetric model with a single $SU(2)_R$ triplet Higgs field [34], which is reproduced if one takes $v_R^{\mathbf{15}} = 0$. This distinguishing feature can be tested at the LHC. Currently, the most stringent bound on m_{Z_R} is given by the resonance searches in Drell-Yan processes [39, 40]. The CMS collaboration gives the most severe limits on this process [40]. Using this analysis⁵, it is found that the present lower bound on the Z_R mass in our model is $m_{Z_R} \gtrsim 3.05$ TeV⁶. We show this limit by the blue shaded area in Fig. 2. We find that m_{Z_R} in our model can

⁴As a simple cross check, we here note that for a generic gauge theory which is spontaneously broken by a VEV of a Higgs field $\langle \varphi \rangle$, the trace of the mass matrix M_{AB}^2 of the gauge bosons is given by

$$\sum_A M_{AA}^2 = 2g_G^2 \langle \varphi \rangle^* T_A T_A \langle \varphi \rangle = 2g_G^2 C_G(R) |\langle \varphi \rangle|^2, \quad (11)$$

where $C_G(R)$ denotes the quadratic Casimir invariant for a representation R of a gauge group G . For example, the vector leptoquark mass $m_{V_{LQ}}^2$ is derived from the relation as

$$6m_{V_{LQ}}^2 = 2g_A^2 C_{SU(4)}(\mathbf{15}) \cdot |\langle (\mathbf{15}, \mathbf{1}, \mathbf{1})_C \rangle|^2. \quad (12)$$

For the TeV-scale gauge bosons, on the other hand, we have

$$\begin{aligned} 2m_{W_R}^2 + m_{Z_R}^2 &= 2g_{2R}^2 C_{SU(2)}(\mathbf{3}) \cdot [|\langle (\mathbf{1}, \mathbf{1}, \mathbf{3}, +2)_C \rangle|^2 + |\langle (\mathbf{1}, \mathbf{1}, \mathbf{3}, 0)_C \rangle|^2] \\ &\quad + 2g_{B-L}^2 \left(2\sqrt{\frac{3}{8}} \right)^2 \cdot |\langle (\mathbf{1}, \mathbf{1}, \mathbf{3}, +2)_C \rangle|^2. \end{aligned} \quad (13)$$

These two relations are consistent with Eqs. (7) and (9).

⁵The CMS limits are given in terms of the two parameters c_u and c_d , which are defined in [41], as $c_{u,d} = (g'^2/2)(g_V^{u,d^2} + g_A^{u,d^2})Br(\ell\ell)$, where $Br(\ell\ell)$ is the branching fraction into lepton pairs, and $g'g_V^{u(d)}$ and $g'g_A^{u(d)}$ are the vector and axial-vector couplings of Z_R with up-type (down-type) quarks, respectively. Using these expressions we find that our model gives $c_u \simeq 2 \times 10^{-3}$ and $c_d \simeq 1 \times 10^{-2}$, which leads to our stated mass bound on Z_R .

⁶ This limit is relaxed slightly if one allows Z_R to decay into right-handed neutrinos.

satisfy this constraint over a wide range of v_R^{15} . The rest of allowed mass range, $3.05 \text{ TeV} \lesssim m_{Z_R} \lesssim 5.2 \text{ TeV}$, can be reached at Run-II of the LHC. For example, the 14 TeV LHC run with an integrated luminosity of 300 fb^{-1} can probe the entire parameter space in Fig. 2 [42]. Finally, we note in passing that the Z_R in our model may explain the dielectron event at 2.9 TeV recently announced by the CMS collaboration [43], although there are inconsistencies with the CMS dilepton bound. These Z_R signatures can possibly be tested in the near future.




Figure 2: m_{Z_R} as a function of v_R^{15} . Here, we set $m_{W_R} = 1.9 \text{ TeV}$. The gray shaded region predicts right-handed neutrino masses to be lighter than the $W_R^\pm$ mass when $y_\Delta < 1$, while the blue shaded area is excluded by the CMS dilepton searches [40].

The VEVs of the bi-doublet Higgs fields Φ_1 and Φ_{15} originally coming from $(\mathbf{1}, \mathbf{2}, \mathbf{2})_C$ and $(\mathbf{15}, \mathbf{2}, \mathbf{2})_C$ of the $\mathbf{10}_C$ and $\mathbf{126}_C$, respectively, induce mixing between the $SU(2)_L$ and $SU(2)_R$ gauge bosons. Through this mixing, $W_R^\pm$ can decay into a W and Z . Therefore, by appropriately choosing the mixing angle, we may explain the diboson excess observed in the ATLAS experiment [10], as we discuss in the next section.

In addition, the VEVs of Φ_1 , Φ_{15} , and $\Delta = (\mathbf{1}, \mathbf{1}, \mathbf{3}, +2)_C$ from the $\mathbf{126}_C$ generate the fermion mass terms via Yukawa interactions. At the TeV scale, the relevant Yukawa terms are given by

$$\begin{aligned} \mathcal{L}_{\text{Yukawa}} = & -\overline{Q}_L \left(y_1^Q \Phi_1 + \tilde{y}_1^Q \tilde{\Phi}_1 \right) Q_R - \frac{1}{\sqrt{24}} \overline{Q}_L \left(y_{15}^Q \Phi_{15} + \tilde{y}_{15}^Q \tilde{\Phi}_{15} \right) Q_R \\ & - \overline{L}_L \left(y_1^L \Phi_1 + \tilde{y}_1^L \tilde{\Phi}_1 \right) L_R + \sqrt{\frac{3}{8}} \overline{L}_L \left(y_{15}^L \Phi_{15} + \tilde{y}_{15}^L \tilde{\Phi}_{15} \right) L_R \\ & - y_\Delta \overline{L}_R^c \Delta L_R + \text{h.c.} , \end{aligned} \quad (14)$$

where c denotes the charge conjugation and $\tilde{\Phi}_{1,15} \equiv \tau_2 \Phi_{1,15}^* \tau_2$. Here, we note that $y_{1,15}^Q = y_{1,15}^L$ and $\tilde{y}_{1,15}^Q = \tilde{y}_{1,15}^L$ hold at the first intermediate scale because of the $SU(4)_C$ symmetry. After the bi-doublet fields acquire VEVs, $\langle \Phi_{1,15} \rangle = \text{diag}(v_u^{1,15}, v_d^{1,15})$, the operators including these VEVs generate a Dirac mass term for the charged SM fields, while the VEV of Δ gives rise to a Majorana mass terms for the right-handed neutrinos. Since both the $\Phi_{1,15}$ and $\Phi_{1,15}^*$ can couple to the SM fermions, with different Yukawa couplings, the form of the Yukawa couplings is similar to that of a generic two-Higgs doublet model⁷. This Yukawa structure in general suffers from large flavor-changing neutral currents [44–46]⁸. In this work, we just assume that the Yukawa couplings are appropriately aligned so that the flavor-changing processes induced by the exchange of the Higgs fields are sufficiently suppressed.

As mentioned above, the lepton Yukawa couplings are equal to the quark Yukawa couplings at the first intermediate scale. As a result, the Dirac Yukawa couplings for neutrinos are sizable. This is potentially problematic since the Majorana masses for right-handed neutrinos in our model are at the TeV scale. To discuss this point, let us investigate the neutrino mass terms. The masses of light neutrinos in our model are given by the ordinary seesaw relation [49],

$$m_\nu \simeq -m_D m_{\nu_R}^{-1} m_D^T, \quad (15)$$

with

$$m_{\nu_R} = y_\Delta v_R^{10}, \quad m_D = (y_1^L v_u^1 + \tilde{y}_1^L v_d^1) - \sqrt{\frac{3}{8}}(y_{15}^L v_u^{15} + \tilde{y}_{15}^L v_d^{15}). \quad (16)$$

The first expression in Eq. (16) induces Majorana masses of $\mathcal{O}(v_R^{10})$. Then, if the lepton Yukawa couplings are as large as the quark Yukawa couplings, m_D becomes as large as quark masses and thus light neutrino masses generically become much larger than $\mathcal{O}(1)$ eV. To get small neutrino masses, in this paper, we suppress m_D by fine-tuning the lepton Yukawa couplings. Notice that this fine-tuning is possible because the VEV of the $(15, 2, 2)$ component breaks both the $SU(4)_C$ and $SU(2)_R$ relations [50]. The breaking of the $SU(4)_C$ relation can be seen in the different factors in front of y_{15}^Q and y_{15}^L , while the breaking of the $SU(2)_R$ relation is realized when $v_u^{15} \neq v_d^{15}$. This allows us to choose the neutrino Dirac Yukawa couplings freely. In this sense, the $(15, 2, 2)$ component is a necessary ingredient to make this model viable⁹.

As we discuss below, we take right-handed neutrino masses to be larger than m_{W_R} to forbid the leptonic decay channels of $W_R^\pm$. On the other hand, the right-handed neutrino

⁷Although there are 4 Higgs doublets below the TeV scale contributing to electroweak symmetry breaking, the $SU(2)_R$ symmetry reduces the number of Yukawa couplings.

⁸However, such a generic flavor structure may explain a 2.4σ excess in the $h \rightarrow \mu\tau$ decay process recently reported by the CMS collaboration [47]. For interpretations of this excess based on the generic two-Higgs doublet model, see Ref. [48].

⁹The same trick can be used to resolve discrepancies among the charged lepton and quark mass relations.

masses are induced by the Yukawa coupling y_Δ . To satisfy this condition with a perturbative y_Δ , we have a lower bound on v_R^{10} , which then implies a lower bound on the Z_R mass, as can be seen from Eq. (7). For instance, for $m_{W_R} = 1.9$ TeV and $y_\Delta < 1$, we have $m_{Z_R} > 2.2$ TeV. The gray shaded region in Fig. 2 shows the area that results in small right-handed neutrino masses. As can be seen from this figure, the LHC constraints on M_{Z_R} are more severe.

Before concluding this section, we briefly discuss constraints on W_R from flavor observables. The presence of W_R in general induces additional flavor violation. Currently, the K_L-K_S mass difference gives the most severe constraint on the mass of W_R , which is roughly given by [51]

$$m_{W_R} \gtrsim \left(\frac{g_{2R}}{g_{2L}} \right) \times 2.5 \text{ TeV} \simeq 1.7 \text{ TeV} , \quad (17)$$

where we have set $g_{2R}/g_{2L} \simeq 0.67$, which is predicted in our model (see Eq. (6)). Hence, a ~ 2 TeV W_R is still allowed by flavor bounds.

3 Diboson signal

The ATLAS collaboration has reported an anomalous excess of events in the search for resonances decaying into a pair of gauge bosons, focussing on the hadronically decaying gauge bosons [10]. These gauge bosons are detected as a fat jet since they come from a heavy resonance and are highly boosted and thus quarks from the gauge boson are reconstructed as a single jet with a large jet radius. By investigating invariant mass distributions constructed from two fat jets, the ATLAS collaboration has observed a narrow resonance around 2 TeV with WZ , WW , and ZZ final states having a local significance of 3.4, 2.6, and 2.9σ , respectively.¹⁰ The tagging selections for each channel are inexact so that sizable events are expected to contaminate other channels. The CMS collaboration also performed a similar search with no discrimination between W and Z bosons, and found a small excess around 1.9 TeV [54]. Recently, the ATLAS collaboration provided an analysis [55] that combines searches for diboson resonances decaying into fully leptonic [56], semi-leptonic [57, 58], and hadronic final states [10]. This still exhibits a 2.5σ deviation around 2 TeV in the WZ channel. The CMS experiment has also presented their semi-leptonic search result and found a slight excess around 1.8 TeV [59], although the collaboration found no deviation for the leptonic decay channel[60]. In light of these results, we here discuss the possibility that $W_R^\pm$ in our model accounts for the excesses observed in the above searches.

The $W_R^\pm$ can decay into $W^\pm$ and Z through mixing with $W^\pm$, and thus can yield diboson signals [11–16]. In addition, it can decay into a pair of quarks as well as $W^\pm h$. If right-handed neutrinos are lighter than the $W_R^\pm$ mass, the $\ell_R^\pm \nu_R$ channels are also open,

¹⁰We note that possible subtleties in this ATLAS analysis have been pointed out in Ref. [52]. For a subsequent study on this fat-jet analysis performed by the ATLAS collaboration, see Ref. [53].

otherwise the $W_R^\pm$ does not have leptonic decay channels. The dijet, third-generation-quark, and $W^\pm h$ resonance searches are also important for determining if $W_R^\pm$ can explain the diboson anomaly. As for the dijet resonance searches, the CMS experiment [61] observed a $\sim 2\sigma$ excess around 1.8 TeV, while the ATLAS collaboration found no significant excess around this region [62]. The CMS collaboration also reported a 2.2σ excess around 1.8 TeV in the $W^\pm h$ channel, where the $W^\pm$ decays leptonically and the Higgs boson h decays to $b\bar{b}$ [63]. However, the ATLAS collaboration did not observe any deviation from the background [64]. The CMS experiment also analyzed the $W^\pm h$ final states in the fully hadronic channel [65] and found no significant excess. Finally, the searches in the top and bottom quark final states exhibit no excess around ~ 2 TeV [66–69].

Considering these results, the authors in Ref. [14] carried out a cross-section fit to the data for a resonance in the vicinity of 1.8–1.9 TeV; the resultant cross sections they obtained are¹¹

$$\sigma(pp \rightarrow W_R \rightarrow WZ) = 5.7_{-3.3}^{+3.6} \text{ fb} , \quad (18)$$

$$\sigma(pp \rightarrow W_R \rightarrow Wh) = 4.5_{-4.0}^{+5.2} \text{ fb} , \quad (19)$$

$$\sigma(pp \rightarrow W_R \rightarrow jj) = 91_{-45}^{+53} \text{ fb} , \quad (20)$$

$$\sigma(pp \rightarrow W_R \rightarrow tb) = 0_{-0}^{+39} \text{ fb} . \quad (21)$$

In our model, the production cross section of $W_R^\pm$ is given as a function of the $W_R^\pm$ mass, m_{W_R} , and the $SU(2)_R$ gauge coupling, g_{2R} . The former is determined by the position of the resonance, while the latter is determined by enforcing gauge coupling unification and is given in Eqs. (6). The partial decay widths of the dijet and tb channels are also determined by these two quantities. On the other hand, the partial decay widths of the WZ and Wh channels depend on the W – W_R mixing angle ϕ_{LR}^W defined by,

$$\begin{pmatrix} W_L \\ W_R \end{pmatrix} = \begin{pmatrix} \cos \phi_{LR}^W & -\sin \phi_{LR}^W \\ \sin \phi_{LR}^W & \cos \phi_{LR}^W \end{pmatrix} \begin{pmatrix} W \\ W' \end{pmatrix} . \quad (22)$$

The size of the mixing angle is expected to be $\phi_{LR}^W = \mathcal{O}(m_W^2/m_{W_R}^2) \sim 10^{-3}$, though it is highly dependent on the Higgs sector of the model. In the minimal left-right symmetric model[34], which is the same as $v_R^{\mathbf{15}} = 0$ in our model, the mixing angle is given by

$$\phi_{LR}^W \simeq \sin 2\beta \left(\frac{g_{2R}}{g_{2L}} \right) \frac{m_W^2}{m_{W_R}^2} , \quad (23)$$

where $\tan \beta$ is the ratio of the VEVs of the bi-doublet Higgs field $(\mathbf{1}, \mathbf{2}, \mathbf{2})_C$. To explain the observed top and bottom quark mass ratio, $\tan \beta \simeq m_t/m_b$ is usually favored in this

¹¹Here, we show the result for $\sigma(pp \rightarrow W_R \rightarrow tb)$ without using the ATLAS $\ell\nu b\bar{b}$ result [67]. As discussed in Ref. [14], this ATLAS search systematically gives limits on the W_R cross sections which are about 2σ smaller than the expected limit for the entire mass range consider. This leads to a strong upper limit on $\sigma(pp \rightarrow W_R \rightarrow tb)$ after the fit of 38 fb which is due to the fitting method used in that work. Considering the possibility of a systematic error in the ATLAS $\ell\nu b\bar{b}$ search with its large effect on the cross section fit shown in Ref. [14], we use the cross sections found when the ATLAS $\ell\nu b\bar{b}$ search is ignored.

model. In our model, on the other hand, both Φ_1 and Φ_{15} contribute to the mixing angle with a similar form to the above expression. Contrary to the minimal left-right symmetric case, the VEVs of these fields can be chosen almost arbitrary thanks to the generic structure of the Yukawa sector; we can always reproduce the quark and lepton masses by appropriately choosing these VEVs and Yukawa couplings. Because of this added freedom in our model, we do not specify the Higgs and Yukawa sectors of our models and regard the mixing angle as just a free parameter of order 10^{-3} . To simplify our discussion, we also assume the right-handed neutrino masses are larger than the W_R mass in order to forbid the $\ell_R^\pm \nu_R$ decay channels. At the end of this section, we briefly discuss the case in which a right-handed neutrino has a mass lighter than m_{W_R} .

The authors in Ref. [14] performed a parameter fitting in an $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$ model to determine the model parameters, g_{2R}/g_{2L} and ϕ_{LR}^W , for which the required cross sections given above are satisfied. Since the W_R sector in our model is the same as that in Ref. [14], we can directly apply their result to our case. According to their 68% CL result,

$$0.5 \lesssim g_{2R}/g_{2L} \lesssim 0.65, \quad 0.0012 \lesssim \sin \phi_{LR}^W \lesssim 0.0016, \quad (24)$$

where they took $m_{W_R} = 1.9$ TeV, the value preferred by the current experimental data. Notice that our model predicts $g_{2R}/g_{2L} \simeq 0.67$ as can be seen in Eqs. (6), and is very close to the preferred region. Therefore, by choosing the mixing angle in the range presented in Eq. (24), our model can explain the ATLAS diboson anomaly without conflicting with other collider bounds.

Using the gauge couplings given in Eq. (6), and a mixing angle in the range found in Eq. (24), we evaluate the production cross section $\sigma(pp \rightarrow W_R^\pm)$, total decay width Γ_{tot} , and the branching ratios of $W_R^\pm$ which are given in the Appendix. The total decay width and branching fractions for our model are summarized in Table. 2. Here, we set $m_{W_R} = 1.9$ TeV. Notice that $\text{BR}(WZ)$ is sizable even though the mixing angle $\sin \phi_{LR}^W$ is very small. This is because the $W_R^\pm \rightarrow W^\pm Z$ decay process is enhanced by a factor of $(m_{W_R}/m_W)^4$ due to the high-energy behavior of the longitudinal mode of $W_R^\pm$. This enhancement compensates for the suppression coming from the mixing angle. In addition, we have $\text{BR}(WZ) \simeq \text{BR}(Wh)$, as a consequence of the equivalence theorem. As for the production cross section, we compute it using MADGRAPH5 [70] at the leading order, and re-scale by the so-called k factor which corresponds to the effects of the higher-order corrections; this value is found to be $k \simeq 1.3$ [71, 72]. The resultant cross section is

$$\sigma(pp \rightarrow W_R^\pm) \simeq 143 \times \left(\frac{g_{2R}}{0.42} \right)^2 \text{ fb}, \quad (25)$$

again for $m_{W_R} = 1.9$ TeV. Considering the fact that about a half of WZ events decay hadronically, and using the acceptance ~ 0.14 [10], we estimate that there are about 9 (5) additional events in total in the presence of $W_R^\pm$ if $\sin \phi_{LR}^W = 0.0016$ (0.0012). This can explain the diboson excess observed in Ref. [10].

As we have seen above, to explain the diboson anomaly, the mixing angle ϕ_{LR}^W should be $\mathcal{O}(10^{-3})$. This size of the W - W_R mixing could be constrained by the electroweak precision

Table 2: Total decay width Γ_{tot} and the branching ratios of $W_R^\pm$ for our model. Here, we set $m_{W_R} = 1.9$ TeV and use the gauge couplings given in Eq. (6).

$\sin \phi_{LR}^W$	Γ_{tot} [GeV]	BR(jj)	BR(tb)	BR(WZ)	BR(Wh)
0.0016	22.3	0.61	0.30	0.047	0.046
0.0012	21.4	0.63	0.31	0.027	0.027

measurements [11, 13]. Indeed, the authors in Ref. [73] derived an upper limit on the mixing angle from the bound on the T -parameter of $|\phi_{LR}^W| \lesssim 5 \times 10^{-4}$ for $m_{W_R} = 2$ TeV, which is smaller than the values in Eq. (24). However, our model also contains Z_R , which affects the electroweak observables as well. Moreover, since its contribution modifies the Z -boson coupling to the SM fermions at tree level through the Z - Z_R mixing, we cannot use the conventional method relying on the S and T parameters [74] to evaluate the electroweak precision constraints. Instead, we need to perform a complete parameter fitting onto the electroweak observables. This is beyond the scope of this paper. We however note that such a parameter fitting has been carried out for the $\text{SU}(2)_L \otimes \text{SU}(2)_R \otimes \text{U}(1)_{B-L}$ models in the literature [71, 75] and, according to the results, a 2 TeV W_R with an $\mathcal{O}(10^{-3})$ W - W_R mixing can evade the electroweak precision constraint. Since the structure of the TeV-scale gauge sector of our model is similar to those in these models, we expect that our model can also avoid this constraint.

Next, we consider the case where a right-handed neutrino has a mass lighter than the W_R mass. This case is potentially interesting since it may explain the 2.8σ anomaly observed by the CMS collaboration in the right-handed neutrino searches looking for dijet plus dilepton events [17]. Indeed, it turns out that W_R with $g_{2R}/g_{2L} \simeq 0.6$ and $m_{W_R} \sim 2$ TeV may explain the excess if the W_R decays into a right-handed neutrino that mainly couples to first generation charged leptons [76], i.e. $W_R^\pm \rightarrow \ell^\pm N_R \rightarrow \ell^\pm \ell^\pm jj$. This simple setup is, however, disfavored once the charge of observed electrons/positrons is considered. The CMS collaboration observed only one same-sign electron event on top of the 13 opposite-sign events. However, if the right-handed neutrino is a Majorana fermion, we expect almost the same numbers of events for each case¹². In addition, such a Majorana right-handed neutrino is severely restricted by the ATLAS search for same-sign leptons plus dijet events [77]. To reconcile these experimental results, there are several possibilities for extending the neutrino sector that work, such as the inverse seesaw [78] or the linear seesaw [79] mechanisms which were discussed in the context of the diboson anomaly in Refs. [15] and [16], respectively. In these cases, heavy neutrinos become pseudo-Dirac fermions, which allows them to evade the bounds from the same-sign leptons plus dijet searches [17, 77]. Although our model potentially accommodates such an extension, we

¹²Since a Majorana fermion and its anti-particle are identical, if a Majorana fermion decays into an electron it would also decay into a positron with the same rate.

leave the investigation of this to future work.

Finally, we discuss future prospects for testing this model. The first step for verifying our model is, of course, to confirm the presence of a ~ 2 TeV W_R at Run-II of the LHC. The production cross section of a W_R in our model at a 13 TeV center-of-mass energy is estimated as ~ 800 fb, where we have used the k factor of $k \simeq 1.2$ [71, 72]. Therefore, a ~ 2 TeV W_R can be tested with an integrated luminosity of ~ 10 fb $^{-1}$. As discussed above, our model may predict a relatively light Z_R , which may also be reached at the LHC. In addition, a (fat) jet plus missing energy search can also probe this model [80], which would constrain the decay mode $WZ \rightarrow \nu\bar{\nu}jj$.

The presence of an $SU(2)_L$ triplet Dirac fermion dark matter is a distinct feature of our model, and thus dark matter searches can also play an important role in testing this model. Since we expect its mass to be around 2 TeV, it is difficult for the LHC to probe it directly. Instead, indirect dark matter detection experiments, such as searches for gamma rays from the Galactic Center or dwarf spheroidal galaxies, are quite promising since this dark matter has a relatively large annihilation cross section. Most work to date on an $SU(2)_L$ triplet fermion dark matter candidate has assumed a Majorana rather than a Dirac particle [81] and there is little work on the Dirac triplet dark matter. To test the dark matter in our model in the indirect detection experiments, it is quite important to evaluate its annihilation cross section precisely and compare it with that of the Majorana dark matter. The direct detection rate of the $SU(2)_L$ triplet dark matter is rather small; its scattering cross section with a nucleon is computed as $\sigma_{SI} \simeq 2 \times 10^{-47}$ cm ^{2} [82]. Still, it is in principle detectable in future experiments, which gives another way to test our model.

4 Conclusion

We have considered a specific (non-supersymmetric) $SO(10)$ model which can simultaneously explain the purported ATLAS diboson excess, provide a stable dark matter candidate, and achieve gauge coupling unification with a suitably high GUT scale. In addition to the gauge and matter sector of the theory which are standard for $SO(10)$, the model has Higgs fields with three different representations under the $SO(10)$ gauge symmetry: a **210** which is used to break $SO(10)$, and plays a role in breaking both of the intermediate scale gauge groups, $SU(4)_C \otimes SU(2)_L \otimes SU(2)_R$ and $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$; a **126** which assist the breaking of the smaller intermediate gauge group and participates in the breaking of the SM gauge symmetries; a **10** which is also involved in the breaking of the SM gauge symmetries. The model also introduces a Dirac **45** which provides us with a stable dark matter candidate in the form of a $SU(2)_L$ triplet. Having fixed the lower intermediate scale at 1.9 TeV (to account for the diboson excess), the GUT scale of order 10^{17} GeV and the high intermediate scale of order 10^9 GeV are determined by requiring gauge coupling unification. In this model, both stages of intermediate symmetry breaking as well as a TeV-scale dark matter triplet are necessary for unification of the gauge couplings.

The high GUT scale ensures proton stability. The relatively high intermediate scale of 10^9 GeV, ensures that the effects induced by the presence of both vector and scalar leptoquarks are small enough to remain compatible with experimental constraints. The lower intermediate scale was chosen to account for the diboson excess, and predicts new right-handed gauge bosons to appear at the TeV scale. In addition to a charged $W_R^\pm$ pair, the model predicts a new Z_R gauge boson whose mass depends on the relative contributions of the two Higgs VEVs participating in the breaking of the low intermediate scale. This set of TeV scale gauge bosons should allow the model to be tested in current and future runs at the LHC.

The contributions of two Higgs VEVs to SM breaking gives the model the flexibility needed to adjust the SM fermion spectrum, thus avoiding some of the overly restrictive predictions often encountered in minimal SO(10) models. This is necessary if one hopes to derive eV scale neutrino masses from the seesaw mechanism given that the Majorana mass for the right-handed neutrino is also at the TeV scale.

Finally, the model contains a testable dark matter candidate and the neutral member of an electroweak triplet. Similar to the often studied supersymmetric wino, the mass splitting is rather small (typically of order 165 MeV), and the annihilation cross section is rather large, requiring the mass of the triplet to be of order 2 TeV as well. Thus the dark matter candidate in this model is testable in indirect dark matter searches as well as direct detection experiments.

Note Added: While completing this work, the CMS collaboration announced a new result for the dijet resonance search using the 13 TeV LHC data with an integrated luminosity of 2.4 fb^{-1} [83]. They give an upper bound on the product of the production cross section (σ), the branching fraction of the dijet channel ($\text{Br}(jj)$), and acceptance (A) of $\sigma A \text{Br}(jj) \lesssim 400 \text{ fb}$ for a resonance mass of 1.9 TeV, and thus we have $\sigma \text{Br}(jj) \lesssim 667 \text{ fb}$ using $A \simeq 0.6$ [83]. The ATLAS collaboration also reported a 13-TeV result for the same channel based on the 3.6 fb^{-1} data, and gave a more severe constraint [84]: $\sigma A \text{Br}(jj) \lesssim 210 \text{ fb}$ for a 1.9 TeV resonance. By using $A \simeq 0.4$ [84], we obtain $\sigma \text{Br}(jj) \lesssim 525 \text{ fb}$. On the other hand, our model predicts $\sigma \text{Br}(jj)$ to be $\sigma \text{Br}(jj) = 421$ (485) fb for a 1.9 TeV W_R with $\sin \phi_{LR}^W = 0.0016$ (0.0012), and thus evades these limits. Since these limits are fairly close to the model predictions, we expect that the presence of W_R in our model can be confirmed in near future.

Acknowledgments

The work of J.L.E., N.N. and K.A.O. was supported by DOE grant de-sc0011842 at the University of Minnesota.

Appendix

A Decay widths

Here, we summarize the partial decay widths of W_R we use in our calculation. For the fermionic channels, $W_R \rightarrow f\bar{f}'$, we use

$$\Gamma(W_R^+ \rightarrow u\bar{d}) = \Gamma(W_R^+ \rightarrow c\bar{s}) = \frac{g_{2R}^2}{16\pi} m_{W_R} , \quad (26)$$

$$\Gamma(W_R^+ \rightarrow t\bar{b}) = \frac{g_{2R}^2}{16\pi} m_{W_R} \left(1 + \frac{m_t^2}{2m_{W_R}^2}\right) \left(1 - \frac{m_t^2}{m_{W_R}^2}\right)^2 . \quad (27)$$

For the $W_R \rightarrow WZ$ decay process, we have

$$\begin{aligned} \Gamma(W_R^+ \rightarrow W^+ Z) &= \frac{g_{2L}^2}{192\pi} \sin^2 \phi_{LR}^W \frac{m_{W_R}^5}{m_W^4} \left(1 - 2 \frac{m_W^2 + m_Z^2}{m_{W_R}^2} + \frac{(m_W^2 - m_Z^2)^2}{m_{W_R}^4}\right)^{\frac{3}{2}} \\ &\times \left(1 + 10 \frac{m_W^2 + m_Z^2}{m_{W_R}^2} + \frac{m_W^4 + 10m_W^2 m_Z^2 + m_Z^4}{m_{W_R}^4}\right) . \end{aligned} \quad (28)$$

Finally, the $W_R \rightarrow Wh$ decay width is given by

$$\begin{aligned} \Gamma(W_R^+ \rightarrow W^+ h) &= \frac{g_{2L}^2}{192\pi} \sin^2 \phi_{LR}^W \frac{m_{W_R}^5}{m_W^4} \left(1 - 2 \frac{m_W^2 + m_h^2}{m_{W_R}^2} + \frac{(m_W^2 - m_h^2)^2}{m_{W_R}^4}\right)^{\frac{1}{2}} \\ &\times \left(1 + \frac{10m_W^2 - 2m_h^2}{m_{W_R}^2} + \frac{(m_W^2 - m_h^2)^2}{m_{W_R}^4}\right) , \end{aligned} \quad (29)$$

in the decoupling limit.

References

- [1] H. Georgi and S. L. Glashow, Phys. Rev. Lett. **32**, 438 (1974).
- [2] H. Georgi, AIP Conf. Proc. **23**, 575 (1975). H. Fritzsch and P. Minkowski, Annals Phys. **93**, 193 (1975).
- [3] M. S. Chanowitz, J. R. Ellis and M. K. Gaillard, Nucl. Phys. B **128**, 506 (1977); H. Georgi and D. V. Nanopoulos, Nucl. Phys. B **155**, 52 (1979);
- [4] H. Georgi and D. V. Nanopoulos, Nucl. Phys. B **159**, 16 (1979); C. E. Vayonakis, Phys. Lett. B **82**, 224 (1979) [Phys. Lett. **83B**, 421 (1979)].
- [5] S. Rajpoot, Phys. Rev. D **22**, 2244 (1980); M. Yasue, Prog. Theor. Phys. **65**, 708 (1981) [Erratum-ibid. **65**, 1480 (1981)]; J. M. Gipson and R. E. Marshak, Phys. Rev. D **31**, 1705 (1985); D. Chang, R. N. Mohapatra, J. Gipson, R. E. Marshak and

- M. K. Parida, Phys. Rev. D **31**, 1718 (1985); N. G. Deshpande, E. Keith and P. B. Pal, Phys. Rev. D **46**, 2261 (1993); N. G. Deshpande, E. Keith and P. B. Pal, Phys. Rev. D **47**, 2892 (1993) [hep-ph/9211232]; S. Bertolini, L. Di Luzio and M. Malinsky, Phys. Rev. D **81**, 035015 (2010) [arXiv:0912.1796 [hep-ph]].
- [6] M. Fukugita and T. Yanagida, In *Fukugita, M. (ed.), Suzuki, A. (ed.): Physics and astrophysics of neutrinos* 1-248. and Kyoto Univ. - YITP-K-1050 (93/12,rec.Feb.94) 248 p. C.
- [7] L. Di Luzio, arXiv:1110.3210 [hep-ph].
- [8] T. Fukuyama, Int. J. Mod. Phys. A **28**, 1330008 (2013) [arXiv:1212.3407 [hep-ph]].
- [9] A. Masiero, Phys. Lett. B **93**, 295 (1980); G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); Q. Shafi, M. Sondermann and C. Wetterich, Phys. Lett. B **92**, 304 (1980); F. del Aguila and L. E. Ibanez, Nucl. Phys. B **177**, 60 (1981); K. S. Babu and S. Khan, Phys. Rev. D **92**, 075018 (2015) [arXiv:1507.06712 [hep-ph]].
- [10] G. Aad *et al.* [ATLAS Collaboration], arXiv:1506.00962 [hep-ex].
- [11] J. Hisano, N. Nagata and Y. Omura, Phys. Rev. D **92**, no. 5, 055001 (2015) [arXiv:1506.03931 [hep-ph]].
- [12] K. Cheung, W. Y. Keung, P. Y. Tseng and T. C. Yuan, Phys. Lett. B **751**, 188 (2015) [arXiv:1506.06064 [hep-ph]]; B. A. Dobrescu and Z. Liu, Phys. Rev. Lett. **115**, no. 21, 211802 (2015) [arXiv:1506.06736 [hep-ph]]; A. Thamm, R. Torre and A. Wulzer, Phys. Rev. Lett. **115**, no. 22, 221802 (2015) [arXiv:1506.08688 [hep-ph]]; Q. H. Cao, B. Yan and D. M. Zhang, Phys. Rev. D **92**, no. 9, 095025 (2015) [arXiv:1507.00268 [hep-ph]]; T. Abe, R. Nagai, S. Okawa and M. Tanabashi, Phys. Rev. D **92**, no. 5, 055016 (2015) [arXiv:1507.01185 [hep-ph]]; B. C. Allanach, B. Gripaios and D. Sutherland, Phys. Rev. D **92**, 055003 (2015) [arXiv:1507.01638 [hep-ph]]; T. Abe, T. Kitahara and M. M. Nojiri, arXiv:1507.01681 [hep-ph]; B. A. Dobrescu and Z. Liu, JHEP **1510**, 118 (2015) [arXiv:1507.01923 [hep-ph]]; A. E. Faraggi and M. Guzzi, Eur. Phys. J. C **75**, no. 11, 537 (2015) [arXiv:1507.07406 [hep-ph]]; P. Coloma, B. A. Dobrescu and J. Lopez-Pavon, arXiv:1508.04129 [hep-ph]; J. H. Collins and W. H. Ng, arXiv:1510.08083 [hep-ph]; B. A. Dobrescu and P. J. Fox, arXiv:1511.02148 [hep-ph]; A. Sajjad, arXiv:1511.02244 [hep-ph]; T. Appelquist, Y. Bai, J. Ingoldby and M. Piai, arXiv:1511.05473 [hep-ph]; K. Das, T. Li, S. Nandi and S. K. Rai, arXiv:1512.00190 [hep-ph]; J. A. Aguilar-Saavedra and F. R. Joaquim, arXiv:1512.00396 [hep-ph]; M. Hirsch, M. E. Krauss, T. Opferkuch, W. Porod and F. Staub, arXiv:1512.00472 [hep-ph].
- [13] Y. Gao, T. Ghosh, K. Sinha and J. H. Yu, Phys. Rev. D **92**, no. 5, 055030 (2015) [arXiv:1506.07511 [hep-ph]].

- [14] J. Brehmer, J. Hewett, J. Kopp, T. Rizzo and J. Tattersall, JHEP **1510**, 182 (2015) [arXiv:1507.00013 [hep-ph]].
- [15] P. S. Bhupal Dev and R. N. Mohapatra, Phys. Rev. Lett. **115**, no. 18, 181803 (2015) [arXiv:1508.02277 [hep-ph]].
- [16] F. F. Deppisch, L. Graf, S. Kulkarni, S. Patra, W. Rodejohann, N. Sahu and U. Sarkar, arXiv:1508.05940 [hep-ph].
- [17] V. Khachatryan *et al.* [CMS Collaboration], Eur. Phys. J. C **74**, no. 11, 3149 (2014) [arXiv:1407.3683 [hep-ex]].
- [18] T. W. B. Kibble, G. Lazarides and Q. Shafi, Phys. Lett. B **113**, 237 (1982).
- [19] L. M. Krauss and F. Wilczek, Phys. Rev. Lett. **62**, 1221 (1989).
- [20] L. E. Ibanez and G. G. Ross, Phys. Lett. B **260**, 291 (1991); L. E. Ibanez and G. G. Ross, Nucl. Phys. B **368**, 3 (1992).
- [21] S. P. Martin, Phys. Rev. D **46**, 2769 (1992) [hep-ph/9207218].
- [22] M. Kadastik, K. Kannike and M. Raidal, Phys. Rev. D **81**, 015002 (2010) [arXiv:0903.2475 [hep-ph]]; M. Kadastik, K. Kannike and M. Raidal, Phys. Rev. D **80**, 085020 (2009) [Erratum-ibid. D **81**, 029903 (2010)] [arXiv:0907.1894 [hep-ph]].
- [23] M. Frigerio and T. Hambye, Phys. Rev. D **81**, 075002 (2010) [arXiv:0912.1545 [hep-ph]]; T. Hambye, PoS IDM **2010**, 098 (2011) [arXiv:1012.4587 [hep-ph]].
- [24] Y. Mambrini, N. Nagata, K. A. Olive, J. Quevillon and J. Zheng, Phys. Rev. D **91**, no. 9, 095010 (2015) [arXiv:1502.06929 [hep-ph]].
- [25] N. Nagata, K. A. Olive and J. Zheng, JHEP **1510**, 193 (2015) [arXiv:1509.00809 [hep-ph]].
- [26] C. Arbelaez, R. Longas, D. Restrepo and O. Zapata, arXiv:1509.06313 [hep-ph].
- [27] S. M. Boucenna, M. B. Krauss and E. Nardi, arXiv:1511.02524 [hep-ph].
- [28] G. R. Farrar and P. Fayet, Phys. Lett. B **76**, 575 (1978); S. Dimopoulos and H. Georgi, Nucl. Phys. B **193**, 150 (1981); S. Weinberg, Phys. Rev. D **26**, 287 (1982); N. Sakai and T. Yanagida, Nucl. Phys. B **197**, 533 (1982); S. Dimopoulos, S. Raby and F. Wilczek, Phys. Lett. B **112**, 133 (1982).
- [29] U. Aydemir, D. Minic, C. Sun and T. Takeuchi, arXiv:1509.01606 [hep-ph].
- [30] T. Bandyopadhyay, B. Brahmachari and A. Raychaudhuri, arXiv:1509.03232 [hep-ph].
- [31] U. Aydemir, arXiv:1512.00568 [hep-ph].

- [32] S. Davidson, D. C. Bailey and B. A. Campbell, Z. Phys. C **61**, 613 (1994) doi:10.1007/BF01552629 [hep-ph/9309310].
- [33] V. A. Kuzmin and M. E. Shaposhnikov, Phys. Lett. B **92**, 115 (1980); T. W. B. Kibble, G. Lazarides and Q. Shafi, Phys. Rev. D **26**, 435 (1982); D. Chang, R. N. Mohapatra and M. K. Parida, Phys. Rev. Lett. **52**, 1072 (1984); D. Chang, R. N. Mohapatra and M. K. Parida, Phys. Rev. D **30**, 1052 (1984); D. Chang, R. N. Mohapatra, J. Gipson, R. E. Marshak and M. K. Parida, Phys. Rev. D **31**, 1718 (1985).
- [34] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974) [Phys. Rev. D **11**, 703 (1975)]; R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 566 (1975); R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 2558 (1975); G. Senjanovic and R. N. Mohapatra, Phys. Rev. D **12**, 1502 (1975).
- [35] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.01589 [astro-ph.CO].
- [36] M. Cirelli, N. Fornengo and A. Strumia, Nucl. Phys. B **753**, 178 (2006) [hep-ph/0512090]; M. Cirelli, A. Strumia and M. Tamburini, Nucl. Phys. B **787**, 152 (2007) [arXiv:0706.4071 [hep-ph]]; M. Cirelli and A. Strumia, New J. Phys. **11**, 105005 (2009) [arXiv:0903.3381 [hep-ph]].
- [37] J. Hisano, S. Matsumoto and M. M. Nojiri, Phys. Rev. Lett. **92**, 031303 (2004) [hep-ph/0307216]; J. Hisano, S. Matsumoto, M. M. Nojiri and O. Saito, Phys. Rev. D **71**, 063528 (2005) [hep-ph/0412403].
- [38] J. Hisano, S. Matsumoto, M. Nagai, O. Saito and M. Senami, Phys. Lett. B **646**, 34 (2007) [hep-ph/0610249].
- [39] G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. D **90**, no. 5, 052005 (2014) [arXiv:1405.4123 [hep-ex]].
- [40] V. Khachatryan *et al.* [CMS Collaboration], JHEP **1504**, 025 (2015) [arXiv:1412.6302 [hep-ex]].
- [41] E. Accomando, A. Belyaev, L. Fedeli, S. F. King and C. Shepherd-Themistocleous, Phys. Rev. D **83**, 075012 (2011) [arXiv:1010.6058 [hep-ph]].
- [42] S. Godfrey and T. Martin, arXiv:1309.1688 [hep-ph].
- [43] CMS Collaboration, CMS-DP-2015-039 (2015).
- [44] J. D. Bjorken and S. Weinberg, Phys. Rev. Lett. **38**, 622 (1977).
- [45] B. McWilliams and L. F. Li, Nucl. Phys. B **179**, 62 (1981).
- [46] O. U. Shanker, Nucl. Phys. B **206**, 253 (1982).

- [47] V. Khachatryan *et al.* [CMS Collaboration], Phys. Lett. B **749**, 337 (2015) [arXiv:1502.07400 [hep-ex]].
- [48] D. Aristizabal Sierra and A. Vicente, Phys. Rev. D **90**, no. 11, 115004 (2014) [arXiv:1409.7690 [hep-ph]]; A. Crivellin, G. D'Ambrosio and J. Heeck, Phys. Rev. Lett. **114**, 151801 (2015) [arXiv:1501.00993 [hep-ph]]; L. de Lima, C. S. Machado, R. D. Matheus and L. A. F. do Prado, arXiv:1501.06923 [hep-ph]; I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik, N. Košnik and I. Nišandžić, JHEP **1506**, 108 (2015) [arXiv:1502.07784 [hep-ph]]; Y. Omura, E. Senaha and K. Tobe, JHEP **1505**, 028 (2015) [arXiv:1502.07824 [hep-ph]]; D. Das and A. Kundu, Phys. Rev. D **92**, no. 1, 015009 (2015) [arXiv:1504.01125 [hep-ph]]; Y. n. Mao and S. h. Zhu, arXiv:1505.07668 [hep-ph]; A. Crivellin, J. Heeck and P. Stoffer, arXiv:1507.07567 [hep-ph]; T. Goto, R. Kitano and S. Mori, Phys. Rev. D **92**, no. 7, 075021 (2015) [arXiv:1507.03234 [hep-ph]]; C. W. Chiang, H. Fukuda, M. Takeuchi and T. T. Yanagida, JHEP **1511**, 057 (2015) [arXiv:1507.04354 [hep-ph]]; A. Crivellin, J. Heeck and P. Stoffer, arXiv:1507.07567 [hep-ph]; F. J. Botella, G. C. Branco, M. Nebot and M. N. Rebelo, arXiv:1508.05101 [hep-ph]; Y. Omura, E. Senaha and K. Tobe, arXiv:1511.08880 [hep-ph].
- [49] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, Conf. Proc. C **7902131**, 95 (1979); M. Gell-Mann, P. Ramond and R. Slansky, Conf. Proc. C **790927**, 315 (1979) [arXiv:1306.4669 [hep-th]]; S. L. Glashow, NATO Sci. Ser. B **59**, 687 (1980); R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980); R. N. Mohapatra and G. Senjanovic, Phys. Rev. D **23**, 165 (1981).
- [50] K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. **70**, 2845 (1993) [hep-ph/9209215]; K. Matsuda, Y. Koide and T. Fukuyama, Phys. Rev. D **64**, 053015 (2001) [hep-ph/0010026]; B. Bajc, A. Melfo, G. Senjanovic and F. Vissani, Phys. Rev. D **73**, 055001 (2006) [hep-ph/0510139]; T. Fukuyama, K. Ichikawa and Y. Mimura, arXiv:1508.07078 [hep-ph].
- [51] Y. Zhang, H. An, X. Ji and R. N. Mohapatra, Nucl. Phys. B **802**, 247 (2008) [arXiv:0712.4218 [hep-ph]]; A. Maiezza, M. Nemevsek, F. Nesti and G. Senjanovic, Phys. Rev. D **82**, 055022 (2010) [arXiv:1005.5160 [hep-ph]]; S. Bertolini, A. Maiezza and F. Nesti, Phys. Rev. D **89**, no. 9, 095028 (2014) [arXiv:1403.7112 [hep-ph]].
- [52] D. Gonçalves, F. Krauss and M. Spannowsky, Phys. Rev. D **92**, no. 5, 053010 (2015) [arXiv:1508.04162 [hep-ph]].
- [53] G. Aad *et al.* [ATLAS Collaboration], arXiv:1510.05821 [hep-ex].
- [54] V. Khachatryan *et al.* [CMS Collaboration], JHEP **1408**, 173 (2014) [arXiv:1405.1994 [hep-ex]].
- [55] ATLAS Collaboration, ATLAS-CONF-2015-045.

- [56] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **737**, 223 (2014) [arXiv:1406.4456 [hep-ex]].
- [57] G. Aad *et al.* [ATLAS Collaboration], Eur. Phys. J. C **75**, no. 2, 69 (2015) [arXiv:1409.6190 [hep-ex]].
- [58] G. Aad *et al.* [ATLAS Collaboration], Eur. Phys. J. C **75**, no. 5, 209 (2015) [Eur. Phys. J. C **75**, 370 (2015)] [arXiv:1503.04677 [hep-ex]].
- [59] V. Khachatryan *et al.* [CMS Collaboration], JHEP **1408**, 174 (2014) [arXiv:1405.3447 [hep-ex]].
- [60] V. Khachatryan *et al.* [CMS Collaboration], Phys. Lett. B **740**, 83 (2015) [arXiv:1407.3476 [hep-ex]].
- [61] V. Khachatryan *et al.* [CMS Collaboration], Phys. Rev. D **91**, no. 5, 052009 (2015) [arXiv:1501.04198 [hep-ex]].
- [62] G. Aad *et al.* [ATLAS Collaboration], Phys. Rev. D **91**, no. 5, 052007 (2015) [arXiv:1407.1376 [hep-ex]].
- [63] CMS Collaboration [CMS Collaboration], CMS-PAS-EXO-14-010.
- [64] G. Aad *et al.* [ATLAS Collaboration], Eur. Phys. J. C **75**, no. 6, 263 (2015) [arXiv:1503.08089 [hep-ex]].
- [65] V. Khachatryan *et al.* [CMS Collaboration], arXiv:1506.01443 [hep-ex].
- [66] G. Aad *et al.* [ATLAS Collaboration], Eur. Phys. J. C **75**, no. 4, 165 (2015) [arXiv:1408.0886 [hep-ex]].
- [67] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **743**, 235 (2015) [arXiv:1410.4103 [hep-ex]].
- [68] S. Chatrchyan *et al.* [CMS Collaboration], JHEP **1405**, 108 (2014) [arXiv:1402.2176 [hep-ex]].
- [69] V. Khachatryan *et al.* [CMS Collaboration], arXiv:1509.06051 [hep-ex].
- [70] J. Alwall *et al.*, JHEP **1407**, 079 (2014) [arXiv:1405.0301 [hep-ph]].
- [71] Q. H. Cao, Z. Li, J. H. Yu and C. P. Yuan, Phys. Rev. D **86**, 095010 (2012) [arXiv:1205.3769 [hep-ph]].
- [72] T. Jezo, M. Klasen, D. R. Lamprea, F. Lyonnet and I. Schienbein, JHEP **1412**, 092 (2014) [arXiv:1410.4692 [hep-ph]].
- [73] C. Grojean, E. Salvioni and R. Torre, JHEP **1107**, 002 (2011) doi:10.1007/JHEP07(2011)002 [arXiv:1103.2761 [hep-ph]].

- [74] M. E. Peskin and T. Takeuchi, Phys. Rev. Lett. **65**, 964 (1990); M. E. Peskin and T. Takeuchi, Phys. Rev. D **46**, 381 (1992).
- [75] K. Hsieh, K. Schmitz, J. H. Yu and C.-P. Yuan, Phys. Rev. D **82**, 035011 (2010) [arXiv:1003.3482 [hep-ph]].
- [76] F. F. Deppisch, T. E. Gonzalo, S. Patra, N. Sahu and U. Sarkar, Phys. Rev. D **90**, no. 5, 053014 (2014) [arXiv:1407.5384 [hep-ph]]; M. Heikinheimo, M. Raidal and C. Spethmann, Eur. Phys. J. C **74**, no. 10, 3107 (2014) [arXiv:1407.6908 [hep-ph]]; F. F. Deppisch, T. E. Gonzalo, S. Patra, N. Sahu and U. Sarkar, Phys. Rev. D **91**, no. 1, 015018 (2015) [arXiv:1410.6427 [hep-ph]].
- [77] G. Aad *et al.* [ATLAS Collaboration], JHEP **1507**, 162 (2015) [arXiv:1506.06020 [hep-ex]].
- [78] R. N. Mohapatra, Phys. Rev. Lett. **56**, 561 (1986); R. N. Mohapatra and J. W. F. Valle, Phys. Rev. D **34**, 1642 (1986).
- [79] S. M. Barr, Phys. Rev. Lett. **92**, 101601 (2004) [hep-ph/0309152]; M. Malinsky, J. C. Romao and J. W. F. Valle, Phys. Rev. Lett. **95**, 161801 (2005) [hep-ph/0506296]; T. Fukuyama, T. Kikuchi and T. Osaka, JCAP **0506**, 005 (2005) [hep-ph/0503201].
- [80] S. P. Liew and S. Shirai, arXiv:1507.08273 [hep-ph].
- [81] T. Cohen, M. Lisanti, A. Pierce and T. R. Slatyer, JCAP **1310**, 061 (2013) [arXiv:1307.4082]; J. Fan and M. Reece, JHEP **1310**, 124 (2013) [arXiv:1307.4400 [hep-ph]]; A. Hryczuk, I. Cholis, R. Iengo, M. Tavakoli and P. Ullio, JCAP **1407**, 031 (2014) [arXiv:1401.6212 [astro-ph.HE]]; B. Bhattacharjee, M. Ibe, K. Ichikawa, S. Matsumoto and K. Nishiyama, JHEP **1407**, 080 (2014) [arXiv:1405.4914 [hep-ph]].
- [82] J. Hisano, K. Ishiwata and N. Nagata, JHEP **1506**, 097 (2015) [arXiv:1504.00915 [hep-ph]].
- [83] V. Khachatryan *et al.* [CMS Collaboration], arXiv:1512.01224 [hep-ex].
- [84] [ATLAS Collaboration], arXiv:1512.01530 [hep-ex].